## **ERRATA**

Magnetic Tuning of the Metal-Insulator Transition for Uncompensated Arsenic-Doped Silicon. W. N. SHAFARMAN, T. G. CASTNER, J. S. BROOKS, K. P. MARTIN, and M. J. NAUGHTON [Phys. Rev. Lett. 56, 980 (1986)].

Equation (5) should read

$$\frac{T_0(N,H)}{T_0(N,0)} \simeq \left[ \frac{1 + \eta H^2/\epsilon}{1 + \eta H^2} \right]^{3\nu(0)} \times \left[ \frac{1 + \eta H^2}{\epsilon + \eta H^2} \right]^{3[\nu(0) - \nu_f]g(H)} .$$
(5)

Spin-Dependent Superelastic Scattering from Pure Angular Momentum States of Na(3P). J. J. Mc-Clelland, M. H. Kelley, and R. J. Celotta [Phys. Rev. Lett. 56, 1362 (1986)].

In Fig. 3(b) (9.26 eV incident energy), the spin asymmetry  $A_{-}$  for  $M_{L}=-1$  atoms (squares) was plotted with the wrong sign, and should be reflected about the x axis. The correct curve goes positive initially, and then becomes negative, going through zero at approximately 30°.

The sentence beginning on line 23 of the first column of page 1365 should read as follows: "At 9.26 eV [Fig. 3(b)], on the other hand,  $A_{-}$  shows a rapid sign change near 30°, going from mostly singlet to almost all triplet domination within 5° of scattering angle."

Sensitivity of a Hopf Bifurcation to Multiplicative Colored Noise. R. Lefever and J. Wm. Turner [Phys. Rev. Lett. 56, 1631 (1986)].

The Fokker-Planck equation appearing [Eq. (3)] as

$$[p(u,\theta,z)]^{-1}\partial_{\tau}p(u,\theta,z) = \cdots + \eta^{-1}(\partial_{z}z + \overline{\sigma}^{2}\partial_{z}z)$$

should read

$$\partial_{\tau} p(u, \theta, z) = \{ \cdots + \eta^{-2} (\partial_{z} z + \overline{\sigma}^{2} \partial_{zz}) \} p(u, \theta, z).$$

The same correction is needed in the Fokker-Planck equation on p. 1633.

Polarization Effects in Exclusive Hadron Scattering. GLENNYS R. FARRAR [Phys. Rev. Lett. 56, 1643]

(1986)].

On p. 1644, second paragraph from bottom of the left-hand side, the two choices used for  $\phi_3$  are (i)  $\phi_3$  and  $\phi_1$  equal and (ii)  $\phi_3 = 0.9\phi_1$ , not  $\phi_3 = -0.9\phi_1$ , as printed.

Measurement of the Angular Correlation between Recoil Velocity and Angular Momentum Vectors in Molecular Photodissociation. G. E. Hall, N. Siva-Kumar, P. L. Houston, and I. Burak [Phys. Rev. Lett. 56, 1671 (1986)].

An erroneous version of Fig. 1 was printed. The correct version is given here.




FIG. 1. Experimental and calculated data for the Q(59)and P(59) lines of CO produced in the photodissociation of OCS and probed by laser-induced fluorescence on the CO  $A^{1}\Pi \leftarrow X^{1}\Sigma$  transition using circularly polarized light. The dissociation and LIF probe lasers were orthogonal. Fluorescence was detected without polarization selection at an angle of 45° to each laser beam. Upper row: Doppler profiles expected in the absence of v-J correlation (solid lines) calculated from Eq. (4); the Gaussian laser linewidth is shown in the dashed curve. Bottom row: Experimental profiles (dots) and profiles calculated for v L J using Eq. (6) (solid curves). Along each row, the left-hand panel is for the Q(59) line with the electric vector of the dissociating light  $\hat{\mathbf{E}}$ aligned perpendicular to the propagation vector of the probe light  $Z(\theta'=90^\circ)$ , the second panel is for the Q(59) line with  $\hat{E} \parallel Z \ (\theta' = 0^{\circ})$ , the third panel is for the P(59) line with  $\hat{E} \perp Z$  ( $\theta' = 90^{\circ}$ ), and the last panel is for the P(59)line with  $\hat{E} \parallel Z \ (\theta' = 0^{\circ})$ . Calculations using either Eq. (4) or (6) are made with values of  $\beta = 0.6$  for the recoil anisotropy, 0.14 cm<sup>-1</sup> for the FWHM laser linewidth, and 1232 m/s for the velocity, as determined by energy and momentum conservation. The relationship between the angular variable X and the detuning is  $\Delta v = (v_0/c)v_0\cos X$ , where  $v_0$ is the center frequency and c is the speed of light.